arXiv:astro-ph/0211009v1 1 Nov 2002

The Rotation Curve and Mass–Distribution in Highly Flattened Galaxies

N. R. Sibgatullin,* A. A. García, V. S. Manko

Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN,
A.P. 14-740, 07000 México D.F., Mexico

Abstract

A new method is developed which permits the reconstruction of the surface–density distribution in the galactic disk of *finite* radius from an *arbitrary* smooth distribution of the angular velocity via two simple quadratures. The existence of upper limits for disk's mass and radius during the analytic continuation of rotation curves into the hidden (non–radiating) part of the disk is demonstrated.

PACS number(s): 98.50

---

*On leave of absence from the Department of Hydrodynamics, Lomonosov Moscow State University, Moscow 119899, Russia

The rotation curves of spiral galaxies supply astronomers with the direct information about distribution of the gravitational potential in galactic disks. The measurements of H_α, HI and CO lines allowed to construct rotation curves for our and nearby galaxies (for early reviews cf. Refs. [1–3]). The advanced modern techniques of observation and data processing permitted to extend rotation curves at the distances far beyond the galactic center and discover the flat rotation curves [3–7]. The first galactic disk models were based on the concept of oblate shells (homoeoids) between confocal spheroids [8–10]. Brandt [11], Brandt and Belton [12] tended the eccentricity of confocal spheroids to zero, thus obtaining the disks covering the whole plane $z = 0$. They proposed a special rotation curve possessing Keplerian asymptotics for M31 and NGC5055, and found numerically the corresponding surface–density distribution. Another approach was developed by Toomre [13] who considered a new class of asymptotically Keplerian rotation curves and found, for infinite disks, the corresponding surface–density distributions. The problem of reconstruction of the surface density from the rotation curve in the case of infinite disks containing a black hole at the center was solved in our work [14]. Mestel [15] pointed out that the highly flattened galaxies and nebulas should be modeled by self–gravitating disks of *finite* radius R. He discussed two limiting cases: a rigidly rotating disk and a model with a constant velocity. In a different context some exact models of self–gravitating disks with outer boundary were found by Morgan and Morgan [16].

In the present paper we propose a new method to the problem of reconstruction of the surface–density distribution in a self–gravitating disk of finite extension from an arbitrary smooth distribution of the angular velocity, and it permits to obtain the *general* solution of this problem in terms of two simple quadratures. We shall also demonstrate that if a rotation curve is analytically continued into the hidden (non–radiating) part of the disk, then the disk's mass can achieve its maximal value at a *finite* radius, and not necessarily at infinity. Like it is done in the classical papers [10–13], we shall assume an idealized model for the highly flattened galactic disks in which the influence of viscosity, pressure, non–disk component and all sort of non–circular motion of matter will be neglected. We will not separate the gravitational fields of the dark (non–radiating) and radiating matter, firstly, because they both exert the same influence on rotation curves, and, secondly, because no reliable mechanism for such separation is yet known [17, 4, 18].

Our approach to the Newtonian self–gravitating disk problem is based on a new representation for the gravitational potential of a disk with the boundary radius R which we take in the form

$$\Phi(\rho, z) = \sqrt{\tilde{\rho}^2 + \tilde{z}^2}\; \tilde{\Phi}(\tilde{\rho}, \tilde{z})\,, \tag{1}$$

where the auxiliary space variables $\tilde{\rho}, \tilde{z}$ are related to the genuine cylindrical coordinates ρ, z via the formulas

$$\tilde{\rho} = R^2\rho/r^2, \quad \tilde{z} = R^2 z/r^2, \quad r^2 \equiv \rho^2 + z^2\,, \tag{2}$$

and the function $\tilde{\Phi}$ is defined in the following way:

$$\tilde{\Phi}(\tilde{\rho}, \tilde{z}) = \frac{1}{2\pi^2}\int\limits_0^\pi \mathrm{d}\theta \int\limits_R^\infty \ln[(s - \tilde{\rho}\cos\theta)^2 + \tilde{z}^2]\, \alpha(s)\, \mathrm{d}s\,, \tag{3}$$

$\alpha(s)$ being a real function which has the meaning of the density of sources distribution. Due to the specific analytical properties of the logarithmic function which turn out to be most advantageous for fulfilling the Laplace equation in the region exterior to disk's sources, hereafter we shall be able to relate the function $\alpha(s)$ to the surface density $\sigma(\rho)$ of the disk and to the distribution of the angular velocity $\omega(\rho)$ via simple integral formulas.

The potential $\Phi(\rho, z)$ which, as can be easily seen from (1), is connected with $\tilde{\Phi}$ by Thomson's transform, satisfies Laplace's equation everywhere except on the disk. On the circle $z = 0$, $\rho \leq R$ the normal derivative $\partial\Phi/\partial z$ experiences a jump. To show this, let us tend z to zero. Then, by virtue of (2), we have

$$\lim_{z\to+0}\frac{\partial\Phi}{\partial z} = \frac{\tilde{\rho}^3}{R^2}\lim_{\tilde{z}\to+0}\frac{\partial\tilde{\Phi}}{\partial\tilde{z}}\,, \tag{4}$$

while from (3) follows that

$$\lim_{\tilde{z}\to+0}\frac{\partial\tilde{\Phi}}{\partial\tilde{z}} = \lim_{\tilde{z}\to+0}\frac{1}{2\pi^2}\int\limits_0^\pi \mathrm{d}\theta \int\limits_R^\infty \left(\frac{-\mathrm{i}}{s - \tilde{\rho}\cos\theta - \mathrm{i}\tilde{z}} + \frac{\mathrm{i}}{s - \tilde{\rho}\cos\theta + \mathrm{i}\tilde{z}}\right)\alpha(s)\, \mathrm{d}s\,. \tag{5}$$

Due to the Sokhotsky–Plemelj formula for the Cauchy-type integrals, we obtain from (5)

$$\left.\frac{\partial\tilde{\Phi}}{\partial\tilde{z}}\right|_{\tilde{z}\to+0} = \begin{cases} 0\,, & 0 \le \tilde{\rho} < R \\ \dfrac{1}{\pi}\displaystyle\int\limits_R^{\tilde{\rho}} \frac{\alpha(s)\mathrm{d}s}{\sqrt{\tilde{\rho}^2 - s^2}}\,, & \tilde{\rho} > R\,, \end{cases} \tag{6}$$

or, returning to the coordinates ρ, z:

$$\left.\frac{\partial\Phi}{\partial z}\right|_{z\to+0} = -2\pi G\sigma(\rho) = \begin{cases} \dfrac{R^4}{\pi\rho^2}\displaystyle\int\limits_R^{R^2/\rho} \frac{\alpha(s)\mathrm{d}s}{\sqrt{R^4 - \rho^2 s^2}}\,, & 0 \le \rho < R \\ 0\,, & \rho > R\,, \end{cases} \tag{7}$$

where $\sigma(\rho)$ is the surface density of the disk.

The condition of balance of the gravitational and centrifugal forces for zero–pressure matter of the disk at $z = 0$, $\rho < R$, i.e.,

$$\left.\frac{\partial\Phi}{\partial\rho}\right|_{z\to 0} + \omega^2\rho = 0\,, \tag{8}$$

can be represented in the form

$$\frac{\partial}{\partial\tilde{\rho}}(\tilde{\rho}\,\tilde{\Phi}) = \frac{\omega^2 R^4}{\tilde{\rho}^3}\,, \tag{9}$$

whence it follows that

$$\tilde{\rho}\frac{\partial\tilde{\Phi}}{\partial\tilde{\rho}} = \frac{R^4}{\tilde{\rho}}\int \tilde{\rho}\left(\frac{\omega^2}{\tilde{\rho}^3}\right)_{,\tilde{\rho}} \mathrm{d}\tilde{\rho}\,. \tag{10}$$

From (3) we obtain the form of the left–hand side of (10) in the limit $z \to 0$:

$$\tilde{\rho}\frac{\partial\tilde{\Phi}}{\partial\tilde{\rho}} = \frac{1}{\pi}\left(\int\limits_R^{\infty}\alpha(s)\,\mathrm{d}s - \int\limits_{\tilde{\rho}}^{\infty}\frac{\alpha(s)s\,\mathrm{d}s}{\sqrt{s^2 - \tilde{\rho}^2}}\right)\,. \tag{11}$$

In the subsequent formulas it is convenient to pass to the non–dimensional variables

$$t \equiv R^2/s^2\,, \quad x \equiv R^2/\tilde{\rho}^2 = \rho^2/R^2\,. \tag{12}$$

In the new variables, formula (7) for the surface–density distribution assumes the form

$$\sigma(x) = -\frac{R}{4\pi^2 G x}\int\limits_x^1 \frac{\alpha(t)\,\mathrm{d}t}{t\sqrt{t - x}} \tag{13}$$

(note that $\sigma(x)$ is not singular at $x = 0$ because of the condition (15) below), so that we are able now to calculate the total mass of the disk:

$$M = 2\pi \int_0^R \sigma(\rho)\rho \, \mathrm{d}\rho = -\frac{R^3}{4\pi G} \int_0^1 \frac{\mathrm{d}x}{x} \int_x^1 \frac{\alpha(t)\,\mathrm{d}t}{t\sqrt{t-x}} \,. \tag{14}$$

For M to be a finite quantity, it is necessary that

$$\lim_{x\to 0} \int_x^1 \frac{\alpha(t)\,\mathrm{d}t}{t\sqrt{t-x}} = \int_0^1 \frac{\alpha(t)\,\mathrm{d}t}{t^{3/2}} = 0 \,. \tag{15}$$

From (14) and (15) a simpler formula for the mass of the disk is readily obtainable:

$$M = -\frac{R^3}{4\pi G} \int_0^1 \frac{\alpha(t)}{t^{3/2}} \ln t \, \mathrm{d}t \,. \tag{16}$$

Taking into account (11) and (15), the balance equation (10) yields the integral equation of the Abel–type with respect to $\alpha(t)$:

$$-\frac{1}{2\pi} \int_0^x \frac{\alpha(t)\,\mathrm{d}t}{t^{3/2}\sqrt{x-t}} = \int_0^x (\omega^2 t^{3/2})'_t \frac{\mathrm{d}t}{\sqrt{t}} + A_0 \,, \tag{17}$$

where the integration constant A_0 should be found from the condition (15).

The solution of equation (17) can be written in the form (cf. Ref. [19], p. 318)

$$\frac{\alpha(x)}{2x^{3/2}} = -\frac{\mathrm{d}}{\mathrm{d}x} \int_0^x \frac{\mathrm{d}t}{\sqrt{x-t}} \Big(A_0 + \int_0^t \frac{1}{\sqrt{s}} (\omega^2 s^{3/2})'_s \, \mathrm{d}s \Big) = -\frac{A_0}{\sqrt{x}} - \int_0^x \frac{(\omega^2 t^{3/2})'_t \, \mathrm{d}t}{\sqrt{t}\sqrt{x-t}} \,. \tag{18}$$

In the general case the surface density has the following asymptotic behavior in the neighborhood of the outer rim $x \to 1$:

$$\sigma(x) \approx A\sqrt{1-x} \,, \quad A = \frac{2R}{\pi^2 G} \int_0^1 \frac{(\omega^2 t^{3/2})'_t \sqrt{t}\,\mathrm{d}t}{\sqrt{1-t}} \,. \tag{19}$$

Formulas (13), (18) provide the *general* solution to the problem under consideration in terms of two successive quadratures: for a given distribution $\omega(\rho)$, one should first find the function $\alpha(\rho)$ with the aid of the formula (18) and then calculate the function $\sigma(\rho)$ from the integral (13). The total mass of the disk is defined by (16).

Let us illustrate the use of the general formulas with some concrete examples. Among numerous classes of smooth rotation curves which can be treated analytically with the aid of our formulas we choose the physically meaningful velocity distributions of the type

$$\omega^2 = \omega_0^2 b^3/(b^2+\rho^2)^{3/2}, \quad \omega_0, b = \text{const.} \tag{20}$$

This particular curve was proposed by Brandt [11] in the case of *infinite* disks and it possesses the Keplerian asymptotics at $\rho \to \infty$; its parameter b multiplied by $\sqrt{2}$ has the meaning of the radius of the circumference at which the linear velocity has its maximal value, while ω_0 is the angular velocity at the center of the disk. A wide family of asymptotically Keplerian rotation curves having the form

$$\omega_n^2 = \frac{\omega_0^2\, b^{2n+2}}{n!}\left(-\frac{\partial}{\partial b^2}\right)^{n-1}\frac{1}{b(b^2+\rho^2)^{3/2}}\,, \quad n = 1, 2, ... \tag{21}$$

and containing (20) as the particular $n = 1$ specialization was studied by Toomre [13]; the corresponding surface–density distributions (in the case of *infinite* disks) can be shown to be given by the formula

$$\sigma_n(\rho) = \frac{\omega_0^2\, b^{2n+2}(2n-1)!!}{2^n n!\,\pi G(b^2+\rho^2)^{(2n+1)/2}}\,. \tag{22}$$

In what follows we shall obtain the analog of the expression (22) in the case of *finite* disks. For this purpose it is advantageous to rewrite the rotation curves (21) in terms of the dimensionless variable $x = \rho^2/R^2$ and parameter $a = b^2/R^2$:

$$\omega_n^2 = \frac{\omega_0^2\, a^{n+1}}{n!}\left(-\frac{\partial}{\partial a}\right)^{n-1}\frac{1}{\sqrt{a}(x+a)^{3/2}} \tag{23}$$

(all these ω_n take the same value at $x = 0$: $\omega_n(0) = \omega_0$).

Let us first consider the case $n = 1$. Setting $n = 1$ in (23) and substituting the resulting ω_1^2 into (18), we obtain

$$\frac{\alpha_1(x)}{2x^{3/2}} = \omega_0^2\left(\frac{\sqrt{x}(3a+x)a}{(a+x)^2} - \frac{2a}{1+a}\right)\,. \tag{24}$$

The corresponding formula for the surface–density distribution is then obtainable from (13), and it has the form

$$\sigma_1(x) = \frac{\omega_0^2\, a^{3/2} b S}{\pi^2 G}\,, \quad S \equiv \frac{\sqrt{(1-x)(a+x)} + (1+a)\mathrm{Arctan}\sqrt{(1-x)/(a+x)}}{(1+a)(a+x)^{3/2}}\,. \tag{25}$$

Due to the linearity of the problem, the surface–density distributions corresponding to the rotation curves (23) will be defined in the general case by the expression

$$\sigma_n(x) = \frac{\omega_0^2\, a^{n+\frac{1}{2}} R}{n!\, \pi^2 G} \left(-\frac{\partial}{\partial a} \right)^{n-1} S\,. \tag{26}$$

In the limit $R \to \infty$, formula (26) reduces to Toomre's expression (22). In Fig. 1 we have plotted the surface–density distribution (25) as the function of ρ/b for different values of the ratio b^2/R^2.

The total mass of the disk defined by the surface density (26) is given by the formula

$$M_n = \frac{2\omega_0^2\, a^{n+1} R^3}{n!\, \pi G} \left(-\frac{\partial}{\partial a} \right)^{n-1} \left(-\frac{1}{1+a} + \frac{1}{\sqrt{a}} \mathrm{Arctan} \frac{1}{\sqrt{a}} \right) , \tag{27}$$

and it can be readily found from (16).

It is worthwhile mentioning that the rotation curves can be also approximated by linear combinations of different functions ω_n^2 defined by (21):

$$\omega^2 = \sum_{n=1}^{N} c_n \omega_n^2\,, \quad \sum_{n=1}^{N} c_n = 1\,, \tag{28}$$

and they will all possess the limit of the rigid–body rotation with a given angular velocity ω_0 near the center and the Keplerian asymptotics at large ρ. The corresponding surface–density distributions and total masses will be given by the formulas

$$\sigma(x) = \sum_{n=1}^{N} c_n \sigma_n(x)\,, \quad M = \sum_{n=1}^{N} c_n M_n(x)\,, \tag{29}$$

and in the neighborhood of the outer rim the surface density will have the following asymptotics:

$$\sigma(x) \approx \frac{\omega_0^2 R \sqrt{1-x}}{\pi^2 G} \sum_{n=1}^{N} c_n \left(\frac{a}{a+1} \right)^n . \tag{30}$$

An interesting important question which we would like to discuss in relation with our formalism is the possible estimation of upper limits of the galactic masses and radii of outer rims since, as is well known, real galactic disks can extend at much larger distances than might be thought from the observations of hot gas [10]. Here we shall restrict our consideration to only the case of analytic continuation of the rotation curves into the region $\rho > R_v$ where R_v is a known radius of the visible part of the galactic disk.

Let an arbitrary smooth distribution $\omega^2(\rho)$ be known from observational data for $0 \leq \rho \leq R_v$. Then the corresponding $\sigma(\rho)$ should be constructed via the formulas (13) and (18) at the supposition that the real radius R of the disk is an unknown parameter to be determined. The upper limit R_{max} for the radius of the disk can be found from the condition $(\mathrm{d}\sigma/\mathrm{d}\rho)_{\rho=R} = 0$ or $A = 0$ (see formula (19)), and the value of R_{max} thus obtained should be subsequently substituted into the formula (16) to get the maximal value of the mass. A different, though equivalent, way of obtaining R_{max} is to find it from the condition of mass' maximum. Hence, for the real radius R and real mass M of the galactic disk we get the inequalities $R_v \leq R \leq R_{max}$ and $M \leq M_{max}$. Mention, that for some rotation curves R_{max} may have finite values, but it can also happen that $R_{max} = \infty$ (like in the case of the pure Toomre's angular–velocity distributions ω_n^2, so that only a linear superposition of different ω_n^2 leads to finite R_{max}).

The existence of maximal values for disks' masses at finite radii for the analytically continued rotation curves constitutes a new physical phenomenon (very similar to the static criteria of stability for the barotropic models of stars [20]) which is absent in infinite disks. For instance, in the case of the angular velocity distribution (28) the condition $A = 0$ yields the algebraic equation

$$\sum_{n=1}^{N} c_n \left(\frac{a}{a+1}\right)^n = 0\,, \tag{31}$$

As can be easily seen, for all $c_n > 0$, $n = \overline{1, N}$, this equation has the unique non–negative root $a = 0$ corresponding to $R = \infty$. But already for $c_1 < 0$, $c_i > 0$, $i = \overline{2, N}$, it will have an additional, positive root $a = a_* > 0$. In the interval $0 < a < a_*$ the surface density turns out to be negative in the vicinity of the outer rim $x = 1$, so expressions (29) are only physically meaningful for $a \geq a_*$, the boundary value of the disk radius being $R_{max} = b/\sqrt{a_*}$.

Let us illustrate the above said with the simplest linear combination of the curves (20) possessing the Keplerian asymptotics:

$$\omega^2 = \frac{4q-1}{3}\omega_1^2 + \frac{4}{3}(1-q)\omega_2^2 = \omega_0^2\left[q\left(\frac{a}{a+x}\right)^{3/2} + (1-q)\left(\frac{a}{a+x}\right)^{5/2}\right] ,$$
$$0 \le q \le 1, \quad 0 < a < 0.5 \,. \tag{32}$$

The corresponding surface–density distribution is given by the formula

$$\sigma(x) = \frac{4q-1}{3}\sigma_1(x) + \frac{4}{3}(1-q)\sigma_2(x) \,, \tag{33}$$

where σ_1 and σ_2 are defined by (25).

The derivative $(\mathrm{d}\sigma/\mathrm{d}\rho)_{\rho=R}$ will be equal to zero when $-1+3a+4q=0$, whence one obtains that a is equal to $(1-4q)/3$.

On the other hand, the mass of the disk with the angular–velocity distribution (32) is equal to

$$M = \frac{2\omega_0^2 b^3}{\pi}\left[\frac{\sqrt{a}(2-5q-3aq)}{3(1+a)^2} + q\,\mathrm{Arctan}\frac{1}{\sqrt{a}}\right] , \tag{34}$$

and, as function of a for fixed $\omega_0^2 b^3$, this expression achieves its maximum at $a_* = (1-4q)/3$.

Therefore, we have arrived at the following important conclusion:

When $0 \le q < 0.25$, the radius of the disk cannot be greater than $b\sqrt{3/(1-4q)}$. For $0.25 \le q \le 1$, as can be seen from (34), the mass of the disk becomes a monotonically decreasing function of a, so that its maximum is achieved in the limit $R \to \infty$. The dependence of M_{max} on q is shown in Fig. 2. This example reveals in particular the difficulties which one may encounter while approximating a real rotation curve by superposing standard rotation curves in the case of infinite disks; these difficulties are absent in the case of disks of finite extension.

This work has been supported by Projects 34222–E and 38495–E from CONACyT of Mexico. N.R.S. acknowledges financial support from SNI–CONACyT, Mexico.

REFERENCES

[1] G. de Vaucouleurs and K.C. Freeman, Vistas in Astronomy **14**, 163 (1972).

[2] E.M. Burbidge and G.R. Burbidge, in *Stars and Stellar Systems IX: Galaxies and the Universe*, edited by A. Sandage, M. Sandage and J. Kristian (University of Chicago Press, 1975), p. 81.

[3] P.C. van der Kruit and R.J. Allen, Ann. Rev. Astron. Astroph. **16**, 103 (1978).

[4] J. Binney and S. Tremaine, Galactic Dynamics (Princeton Univ. Press, 1987).

[5] J. Binney and M. Merrifield, Galactic Astronomy (Princeton Univ. Press, 1998).

[6] Y. Sofue and V.C. Rubin, Ann. Rev. Astron. Astroph. **39**, 137 (2001).

[7] M. Persic, P. Salucci and F. Stel, M.N.R.A.S. **281**, 27 (1996).

[8] H.W. Babcock, Lick. Obs. Bull. **19**, 41 (1939).

[9] E.M. Burbidge, G.R. Burbidge and K.H. Prendergast, Astrophys. J. **130**, 739 (1959); ib. **131**, 282 (1960); ib. **132**, 640 (1960).

[10] M. Schmidt, Bull. Astron. Inst. Neth. **14**, 17 (1957).

[11] J.C. Brandt, Astrophys. J. **131**, 293 (1960).

[12] J.C. Brandt and M.J.S. Belton, Astrophys. J. **136**, 352 (1962).

[13] A. Toomre, Astrophys. J. **138**, 385 (1963).

[14] N.R. Sibgatullin, A.A. García and V.S. Manko, astro–ph/0209200 (accepted for publication in Astronomy Lett.)

[15] L. Mestel, M.N.R.A.S. **126**, 553 (1963).

[16] T. Morgan and L. Morgan, Phys. Rev. **183**, 1097 (1969).

[17] D. Burstein and V.C. Rubin, Astrophys. J. **297**, 423 (1985).

[18] M. Persic and P. Salucci, M.N.R.A.S. **234**, 131 (1988).

[19] I.N. Sneddon, The Use of Integral Transforms (New York: McGraw–Hill Book Company, 1956).

[20] J.L. Tassoul, Theory of Rotating Stars (Princeton: Princeton Univ. Press, 1978).

FIGURES



FIG. 1. The form of the surface density distribution (25) for several particular values of a ($a = 0, 0.16, 0.25, 0.35, 0.5$).



FIG. 2. The dependence of the maximally possible mass of the disk on the parameter q in the example given by formulas (32)–(34).